

Algal acute toxicity of Silver Cyanide in freshwater environment

Jelle Mertens¹ and Tina Liesirova¹

INTRODUCTION

- Silver ions (**Ag⁺**) are known for their natural antimicrobial properties and have a recognised high ecotoxic potential.
- Silver nitrate** is commonly used as test item for Ag⁺ in ecotoxicity tests.
- Algae** are identified as most sensitive aquatic species towards Ag⁺ toxicity (Arijs *et al.* 2021).
- Silver-cyanide complexes** are assumed to be stable at ecologically relevant conditions (Xue *et al.* 1985), but are currently void of reliable ecotoxicity test data.
- Aim of this work: investigate the comparative toxicity of silver nitrate vs. silver-cyanide complexes towards algae.**

INITIAL CONSIDERATIONS

- Two industrially relevant silver-cyanide complexes registered under EU-REACH: **silver cyanide [AgCN]** and **potassium dicyanoargentate [KAg(CN)₂]**.
- Water solubility: AgCN 1.1 µg/L vs. KAg(CN)₂ 200 g/L.
- Speciation calculations (MINEQL) in simple aquatic systems (fixed pH of 5.5, 7 or 8.5; total added Ag concentration 10⁻³ – 10⁻⁹ M) suggest **dominance of Ag-cyanide complexes over Ag⁺ at ecotoxicologically relevant Ag-cyanide concentrations.**




Figure 1: Modeled distribution (using MINEQL) of Ag⁺ (yellow), AgCN (white) and Ag(CN)₂ (black) at pH 5.5 (triangles), 7 (diamonds) and 8.5 (circles) in water after addition of AgCN (left figure) or KAg(CN)₂ (right figure) at concentrations of 1x10⁻⁹ to 1x10⁻³ M

DOSE-RANGE FINDING STUDY

- AgNO₃** and **KAg(CN)₂** tested at 0,5 – 50 µg Ag_{total}/L, **AgCN** tested at 25 – 2500 Ag_{total}/L
- Measured dissolved Ag (% of total added Ag): 1.1-1.3% for AgCN, 47-100% for KAg(CN)₂ and 20-27% for AgNO₃
- Experimental data covering ±25-75% reduction in algal growth (*Raphidocelis subcapitata*)
- Indicative Ag concentrations at 50% growth inhibition [EC50 value]:**
AgNO₃ (0.55 µg Ag/L) < AgCN = KAg(CN)₂ (2.6 - 3.4 µg Ag/L)




Figure 2: Results of the dose-range finding study with AgNO₃ (grey), KAg(CN)₂ (yellow) and AgCN (black). Algal growth (% of control) as a function of measured dissolved Ag concentrations.

MAIN ALGAL TOXICITY STUDY

- AgCN** as test item; added at 6 test concentrations (0.005 - 15 mg Ag_{total} /L)
- Algal toxicity test (*R. subcapitata*) according to OECD guideline 201; GLP compliant
- 72 hour-growth inhibition as endpoint, dissolved Ag measured via ICP-MS
- EC50 value: 16.2 µg Ag/L** [95% confidence interval: 11.3 – 23.1]




Figure 3: Results of the main study with AgCN. Algal growth (% of control) as a function of measured dissolved Ag concentrations.

CONCLUSION

- Speciation modelling suggests stability of silver-cyanide complexes at ecotoxicologically relevant concentrations.
- Results from a dose-range finding study with *Raphidocelis subcapitata* confirm
 - a comparable toxicity of AgCN and KAg(CN)₂
 - a lower toxicity of AgCN and KAg(CN)₂ compared to silver nitrate.
- The toxicity of AgCN in a guideline conform test with *R. subcapitata* results in an EC50 value of 16.2 µg Ag/L.
- The toxicity of silver-cyanide complexes is more than 15-fold lower than of silver nitrate, based on EC50 values (EC50 silver nitrate 0.96 µg Ag/L; Schlich 2016).
- Next step, in this research program, is the determination of the ecotoxicity of silver-cyanide complexes towards invertebrates and fish.

¹ European Precious Metals Federation (EPMF), Brussels, Belgium - E-mail contact: jelle.mertens@epmf.be